

# Chemistry in Action: Space Shuttle Fuel Chemistry



**Ghanshyam L. Vaghjiani**

***Space and Missile Propulsion Division  
Propellant Branch***

***Air Force Research Laboratory  
AFRL/PRSP***

***10 E Saturn Blvd  
Edwards AFB, CA 93524, USA***

***Tel: 661 275 5657***

***Fax: 661 275 6245***

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>AUG 2005</b>		2. REPORT TYPE		3. DATES COVERED -	
4. TITLE AND SUBTITLE <b>Chemistry in Action: Space Shuttle Fuel Chemistry (Briefing Charts)</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) <b>Ghanshyam Vaghjiani</b>				5d. PROJECT NUMBER <b>2303</b>	
				5e. TASK NUMBER <b>0423</b>	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Air Force Research Laboratory (AFMC),AFRL/PRSP,10 E. Saturn Blvd.,Edwards AFB,CA,93524-7680</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>N/A</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES <b>31</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			



# Outline

- **Student's Perception of Chemistry**
- **Role In Science and Technology**
  - **Traditional Areas**
  - **Recent and Emerging Technologies**
- **Space Shuttle-Atmospheric Interactions**
- **New Hypergolic Fuels**
- **Closing Remarks**
  - **Acknowledgements**
  - **Career in the Government**
  - **Web Resources**



# Student's Perception of Chemistry

- ☐ It is too Hard! Too Much Math! I do not Like Cooking!
- ☐ It is Only for Academicians!
- ☐ What use is it for Getting Good Jobs?
- ☐ I Also Thought This! Until I met my Mentor, Ian Worthington

## ☐ Definition:


- ☐ Study of MATTER and the Changes That Take Place With That MATTER

## ☐ Importance:

- ☐ MATTER is Everywhere! Therefore it Matters a lot!
  - ☐ To Understand the Energetics of Breaking and Making Chemical Bonds
  - ☐ We Seek Microscopic Explanation of Macroscopic Changes we Experience



# Role in Science and Technology





# Space Shuttle Propulsion System



## ☐ Space Propulsion (PRC, OMS, Veneers):

### ☐ Hypergolic Liquids



☐ NO External Ignition Required!

## ☐ Boost Phase (2 x 3.1 Mlb):


### ☐ Solids



☐ One-time Squib

## ☐ Launch (3 x 0.4 Mlb):

### ☐ Cryogenic Liquids



☐ One-time Torch





# Space Shuttle-Atmospheric Interactions

## AFRL's Motivation:

- Understand Chemiluminescent Processes at  $\geq 200$  Km

## Strong Emissions From CO(a):




## Cause of Chemiluminescence:

- **Rocket Plume-Atmospheric Interactions**

## UV-Chemistry Questions:

- Precursors?
- Its Formation?
- Its Reactions?

## Space Experiment



### Observation Platforms

Space Shuttle  
Mir Space Station  
MSX

### Thrusters

Space Shuttle  
Progress-M  
Soyuz-TM



# Proposed CO(a) Source Chemistry


■ Unreacted  $\text{CH}_3\text{NHNH}_2 \rightarrow \rightarrow \text{Precursor(s)}$

■ Precursor(s) + O  $\rightarrow$  Products




□ 200 km-Thermosphere

□  $[\text{O}] \gg [\text{O}_2]$




{  
CO + H<sub>2</sub> (Main)  
2H + CO (~ 20%)  
H + HCO (HCO\*)  
CH + OH (~ 6%)



{  
CO + H (Main)  
HCO (HCO\*)  
HCO<sup>+</sup> + e<sup>-</sup> (~ 0.03%)  
C + OH (? +ve E<sub>a</sub>)



# Apparatus








# CO(A) Source Reactions

■ Chemiluminescence Intensity Varied as (Laser Fluence)<sup>2</sup>




■ Diatomics or Triatomics Need to be Internally Excited




# Comparison of CO & OH-Chemiluminescence

Strong CO(A) Signal in O/O<sub>2</sub>




Very Weak CO(A) Signal in O<sub>2</sub> only

Weakened OH(A) Signal in O/O<sub>2</sub>



Strong OH(A) Signal in O<sub>2</sub> only



CO + OH (~ 20%)  
CO<sub>2</sub> + H (~ 30%)  
HCO + O (~ 20%)  
H + CO + O (~ 30%)  
CO + **OH(A)** (~ 0.48%)



# Time-Resolved CO(A)-Chemiluminescence

## ☐ Bimolecular Reaction Rate Coefficients of Added Substrate When CH<sub>4</sub> Present



$$k_{O_2} = (2.2 \pm 0.3) \times 10^{-11}$$

$$k_{N_2O} < 7 \times 10^{-14}$$


$$k_{NO} = (3.4 \pm 0.5) \times 10^{-11}$$

$$k_{H_2} < 2 \times 10^{-13}$$

$$k_{CH_4} < 6 \times 10^{-14}$$




## ☐ (C + O) not the Source





# $\text{CHBr}_3$ Versus $\text{CBr}_4$ Photolysis



☐ Stronger VUV Signal in  $\text{CHBr}_3$  Photolysis



$(\text{CH}^\# \text{ (or } \text{CHBr}^\#) + \text{O})$  Important


☐ Signal in  $\text{CBr}_4$  Photolysis Varies as  $(\text{Fluence})^2$



$(\text{CBr}_2^\# + \text{O})$  not Important, Since  $\text{Br}_2^*$  Signal Varies as  $(\text{Fluence})^1$



# $\text{CBr}_4$ Photolysis



☐  $\text{CBr}_2$  Formed in  
Absence of Photolysis


☐  $\text{CBr}_2$  Formed in  
Photolysis



$\text{CBr}_2 + \text{O} \rightarrow \text{CO}^* + \text{Br}_2$   
not Important



# CHBr<sub>3</sub> Versus CBr<sub>4</sub> Photolysis



□ CHBr<sub>3</sub>

$$k_{O_2} = (2.2 \pm 0.3) \times 10^{-11}$$

□ CBr<sub>4</sub>

$$k_{O_2} = (2.4 \pm 0.4) \times 10^{-12}$$



(CBr<sup>#</sup> + O) Source is not as Important as (CH<sup>#</sup> + O) in CHBr<sub>3</sub> Photolysis

□ CHBr<sup>#</sup> has Very Short Lifetime (~ 5 μs) and  $k_{(CHBr + O_2)} < 2 \times 10^{-14}$




(CHBr<sup>#</sup> + O) Source not Important in CHBr<sub>3</sub> Photolysis



# CH(a<sup>4</sup>Σ<sup>-</sup>) + O

## Reaction Rate Coefficient

Excess CH<sub>4</sub>, 2 Torr He



□  $k_{(CH(a) + O)} = (1.35 \pm 0.47) \times 10^{-10}$

Previously:

□  $k_{(CH(X) + O)} = (9.5 \pm 1.4) \times 10^{-11}$



# Space Shuttle-Atmospheric Interaction: Conclusions

## ● 248-nm Photolysis of $\text{CHBr}_3$ /O-atom Mixtures

### Strong Emissions From:

- $\text{CO(A)}$ ,  $\text{CO(a)}$
- $\text{OH(A)}$  when  $\text{O}_2$  Present
- $\text{Br}_2(\text{D})$

### Kinetic & Laser Fluence Trend Analyses of the Chemiluminescence:

- $\text{CH}(\text{X}^2\Pi, \text{a}^4\Sigma^-) + \text{O}$
- $\text{CBr}_2 + \text{O}$

● Plume Fragments (CH) + Thermosphere (O-atoms) → UV Emissions



# New Hypergolic Fuels

## ✚ AFRL's Motivation:

- Replace Highly Toxic  $\text{CH}_3\text{NHNH}_2$  (MMH)
- Design Better Performing Fuels

## ✚ AFRL's Approach:

- Tune Fuel Structure for;
  - Energy Content: High Heat of Combustion
  - Oxygen Balance: Lower Spacecraft Mass
  - Physical Properties: Higher  $\rho$ , Lower  $m_p$ ,  
Reduced Sensitivities
- Ignition/Combustion Behavior: Short ID Time



Scape Suit

## ✚ Propellant Performance ( $I_{sp}$ )

Fuel + Oxidizer  $\rightarrow$  Products +  $\Delta H$

$$\Delta H = \text{K.E} = \frac{1}{2}mv^2$$

$$I_{sp} = (1/g) \int F(t)dt / \int \dot{M}(t)dt = (1/g)(2\Delta H/m)^{1/2}$$

Cost Reduction in Launch/Health/  
Environment



Splash Shield



# Search For Hypergolic Fuels

## ❑ **CEA-Evaluation:** Identify Better Fuels

	N <sub>2</sub> O <sub>4</sub> /MMH	N <sub>2</sub> O <sub>4</sub> /HEHN	N <sub>2</sub> O <sub>4</sub> /HEATN
KE(MJ kg <sup>-1</sup> )	4.7	3.9	4.0
ρ(kg m <sup>-3</sup> )	1189	1424	1454
FOM	1.0	1.03	1.05

❑ **Definition:** A Pair of Compounds, Upon Contact, Chemically React and Release Sufficient Heat to Spontaneously Ignite

❑ **Discovery/Research of Hypergolic Propellants:** 1930's, Germany (e.g. BMW)

❑ **No *a Priori* Method to Predict Hypergolicity:** **NEW** Fuel & Oxidizer Hypergol Pair Must be Experimentally Verified!



# Screening Fuels For Hypergolicity



Drop-test Apparatus Employed: O/F = ~ 20


Fuel	IRFNA	N <sub>2</sub> O <sub>4</sub>	WFNA
CH <sub>3</sub> NHNH <sub>2</sub> (L) (MMH)	HGI	HGI	HGI
HOCH <sub>2</sub> CH <sub>2</sub> N <sup>+</sup> H <sub>2</sub> NH <sub>2</sub> NO <sub>3</sub> <sup>-</sup> (L) (HEHN)	HGI*	VR	HGI*
(1-ethan-2-ol)-4-amino-1,2,4-triazolium nitrate (L) (HEATN)	SR	VR	
1H-1,2,3-triazole (L)	SR	SR	
1-amino-1,2,3-triazole (M)	HGI*		
3-methyl-1-amino-1,2,3-triazolium nitrate (S)	VR	VR	
▽≡H (L)	VR	VR	VR
▽≡▽ (L)	HGI*	HGI*	HGI*
▽≡≡▽ (L)	HGI*	HGI*	HGI*

HGI=hypergolic ignition, VR=vigorous reaction, SR=slow reaction. At room temperature, fuel is solid (S), liquid (L), or heated to its melting point (M)

\*New hypergols



# Fuel Functionality Affects Ignition




WFNA /  $\nabla \equiv \nabla$  is Hypergolic

$\nabla \equiv \text{H}$

Not Hypergolic

$\nabla \equiv \equiv \nabla$

Is Hypergolic;  
ID = 5.0 ms





# Complexity of the Pre-ignition Chemistry



$\nabla \equiv \nabla / \text{N}_2\text{O}_4$   
ID = 40.6 ms

Unregistered





# New Hypergolic Fuels: Conclusions

## ☐ Characterization of Pre-ignition Chemistry is the Key for Designing new Hypergols

### ☐ Apply Spectroscopic Probing Tools

- ☐ Rapid-Scan FTIR
- ☐ Time-Resolved Raman
- ☐ Time-Resolved Emission
- ☐ High Speed Video

### ☐ Develop Global Initiatory Mechanism

### ☐ Construct Pre-Ignition Models

### ☐ Kinetic Modeling of Ignition

### ☐ Tune Fuel Chemical Functionalities

### ☐ Apply Quantum Chemistry Tools

- ☐  $\Delta H$  of Intermediates
- ☐ PES (Reaction Coordinates)
- ☐ Reaction Rates

### ☐ Provide Initial Rationale to Experimental Observations

**Focused/Intelligent  
Approach to new  
Synthesis of Hypergolic  
Fuels**



# Closing Remarks

## ● Acknowledgements:

### ☐ AFOSR

■ Drs. M. Berkin & M. Berman (\$\$\$\$)

### ☐ AFRL/PRSP

■ Drs. Alfano (**Experimental**), Mills & Boats (**Theory**), Suri & Hawkins (**Synthesis**)

## ● Career in the Government:

### ☐ DoD

■ AFRL, ONR, ARL, etc

### ☐ DoE

■ LLNL, ANL, ONL, LANL, etc

### ☐ DoC

■ NOAA, NIST, etc

### ☐ NASA

■ Dryden, Ames, JPL, etc

### ☐ And Many More .....

## ● Web Resources:

### ☐ American Chemical Society

[www.chemistry.org](http://www.chemistry.org)

### ☐ Edwards AFB

[www.edwards.af.mil](http://www.edwards.af.mil)

### ☐ NASA

[www.nasa.gov](http://www.nasa.gov)

### ☐ New Scientist

[www.sciencesjob.com](http://www.sciencesjob.com)




# Backup Slides





# UV/Vis Plumes

## Radiance Data



⇔ **Plume Data** ⇔



**Modeling Studies**



**Laboratory Studies**




**Chemiluminescent Processes**




**Identify Spacecraft Atmospheric Interactions**

## Spectral Data





# 282.2-nm Signal



## □ Absence of O-atoms

X-trace: (O<sub>2</sub>, 8.8 x 10<sup>14</sup>)

Δ-trace: (O<sub>2</sub>) + (CH<sub>4</sub>, 5.0 x 10<sup>15</sup>)



## □ 5.0 x 10<sup>13</sup> of O-atoms

■-trace: (O<sub>2</sub>, 8.8 x 10<sup>14</sup>)


□-trace: (O<sub>2</sub>) + (CH<sub>4</sub>, 5.0 x 10<sup>15</sup>)




(CBr<sub>2</sub> + CH<sub>4</sub>) Slow Reaction



# $\text{Br}_2^*$ -Chemiluminescence




☐ **Laser off**



☐ **Laser on**


$$\text{Br}_2^* \propto (\text{Fluence})^1$$





# Time Resolved $\text{Br}_2^*$ -Signal

289.9-nm chemiluminescence signal (counts)



Fast  $\text{Br}_2^*$  Rise

Also:

$$k_{\text{O}_2} < 9 \times 10^{-14}$$

$$k_{\text{CH}_4} < 7 \times 10^{-14}$$


$$k_{\text{O}} = (5.4 \pm 1.0) \times 10^{-11}$$



Less Important




Since:





# CO\* Production Mechanism





# Hypergolic Action

- ❑ **No *a Priori* Method:** Hypergolicity Between any Pair of Fuel & Oxidant System Must be Experimentally Verified



- ❑ **Know Your Calories:** < 0.05 cc of a Fuel can Lead to a Spectacular Interaction With an Oxidizer

